

Attention Deficit Hyperactivity Disorder and Dysregulated Eating

Janice Arsenault BSc, RN, MN-NP¹ and Kathryn Weaver BN, RN, MN, PhD²

¹*ADHD and Beyond, New Brunswick, Canada*

²*University of New Brunswick, Canada*

¹*arsenaultnp@gmail.com, ²kweaver@unb.ca*

Abstract

This study was conducted to identify features of dysregulated eating in the context of children living with Attention Deficit Hyperactivity Disorder (ADHD). A deeper understanding of the association between various eating behaviors and specific moderators unique to ADHD was sought to help healthcare providers identify measurable and predictive features that direct nutritional assessment, support, and interventions. Using the Fineout-Overholt, Melnyk, Stillwell, and Williamson (2010) method, seven studies that included children aged 4 and 15 from the Czech Republic, Greece, Germany, Iran, and Korea were reviewed. Results revealed dysregulated eating patterns involving non-traditional eating schedules, increased eating frequency, episodes of overeating, preference for calorically dense food, high intake of sugary foods and beverages, and diets of higher amounts of processed and fast foods. In conclusion, underlying symptomology characteristics such as impulsivity, inattention, and emotional dysregulation may serve to mediate dysregulated eating behaviors in children living with ADHD. Early recognition and intervention could alter the progression of dysregulated eating behavior and its negative sequelae.

Keywords: ADHD, Attention deficit disorder with hyperactivity, Child, Eating behavior

1. Introduction

Inattention, hyperactivity, and impulsivity are the primary symptoms of Attention Deficit Hyperactivity Disorder (ADHD). These symptoms begin in childhood and can be mild, moderate, or severe as they continue into adulthood [1][2]. Attention deficit hyperactivity disorder is subdivided into three subtypes based on symptom presentation: (a) predominantly inattentive or easily distracted and unable to stay on task, (b) predominantly hyperactive-impulsive or being on the go, unable to wait their turn, and reacting before considering consequences, and (c) a combined type defined by both inattentive and hyperactive symptoms [2]. The symptoms of ADHD adversely affect academic, behavioral, and social-emotional functioning [3]. To illustrate, approximately 20-30% of children living with ADHD meet the criteria for depression and experience a 5.5 times greater risk of developing a depressive disorder than adolescents without ADHD [4].

It has been noted that deficits in self-regulation are expected in a substantial number of children living with ADHD and that eating dysregulation may be an early indicator of ADHD [5]. Available research findings suggest that the lifetime prevalence of ADHD is significantly associated with the lifetime prevalence of eating disorders [6][7][8][8]. The most prevalent is

Article history:

Received (August 4, 2022), Review Result (October 5, 2022), Accepted (December 15, 2022)

Binge Eating Disorder (BED), characterized by episodes of eating large amounts of food that are more than what most individuals would eat in a similar period [10]. The impact of dysregulated eating is significant during childhood as this is a critical period of growth and development. In effect, children experiencing ADHD and dysregulated eating patterns were found to be at 1.5 times increased risk for being overweight compared to age-matched controls [7][8][8][11]. Negative outcomes may include obesity, slow growth, shorter stature, suboptimal cognitive development, and psychosocial complications (e.g., low self-esteem, poor body image, depression), both concurrently and long-term [12][13]. Health risks associated with obesity include hypertension, cardiovascular disease, and type-2 diabetes [13][14][15].

Research has focused on adults between the ages of 18 and 44 or adolescents with diagnosed eating disorders as defined in [Table 1] [16][17]. To date, the specific eating characteristics and patterns in children with Attention deficit hyperactivity disorder (ADHD) are not well understood. Exploring the association between various eating behaviors and specific moderators unique to ADHD could have clinical practice significance, such as the importance of healthcare providers screening children who present with ADHD symptomatology for increased risk of disordered eating and associated health problems. Early recognition and intervention could help alter the development or maintenance of dysregulated eating behavior and its negative sequelae. Accordingly, this study was conducted to evaluate the association between core characteristics of ADHD and distinct dysregulated eating patterns and behaviors.

Table 1. Overview of criteria for categories of eating disorders

Anorexia Nervosa (AN) Diagnosis	Related Otherwise specified feeding or eating disorder (OSFED)
<p>All of the following criteria for AN are met:</p> <p>A. Food restriction + low body weight relative to peers</p> <p>B. Intense fear of gaining weight</p> <p>C. Distorted body image.</p> <p>Mild: Body Mass Index (BMI) ≥ 17 kg/ m²</p> <p>Moderate: BMI 16-16.99</p> <p>Severe: BMI 15-15.99</p> <p>Extreme: BMI < 15</p>	<p>Atypical AN: All criteria for AN are met, except that despite significant weight loss, the individual's weight is within or above the normal range.</p>
<p>AN Subtypes:</p> <p>Restricting type: Weight loss is accomplished primarily through dieting, fasting, and excessive exercise. Has not engaged in recurrent episodes of binge eating or purging behavior (i.e., self-induced vomiting or the misuse of laxatives, diuretics, or enemas) during the last three months.</p> <p>Binge-eating/purging type: Has engaged in recurrent episodes of binge eating or purging behavior (i.e., self-induced vomiting or the misuse of laxatives, diuretics, or enemas) during the last three months,</p>	
Bulimia Nervosa (BN) Diagnosis	Related OSFED
<p>All BN criteria met:</p> <p>A. Recurrent episodes of binge eating (eating within a discrete period, an amount of food more significant than what most individuals would eat in a similar period under a similar context) with a concurring sense of lack of control over eating</p> <p>B. Recurrent inappropriate compensatory behaviors to prevent weight gain (e.g., self-induced vomiting; misuse of laxatives,</p>	<p>BN (of low frequency and limited duration): All criteria for BN are met, except that binge eating and inappropriate compensatory behaviors occur, on average, less than once a week and for less than three months.</p>

<p>diuretics, or other medications; fasting; or excessive exercise).</p> <p>C. The binge eating and compensatory behaviors both occur, on average, at least once a week for three months.</p> <p>D. Self-evaluation is unduly influenced by body shape and weight.</p> <p>E. The disorder does not occur exclusively during episodes of AN.</p> <p>Severity: Mild: 1-3 episodes of compensatory behaviors /week Moderate: 4-7 per week Severe: 8-13 per week Extreme: ≥14 per week</p>	
<p>Binge Eating Disorder (BED) Diagnosis</p>	<p>Related OSFED</p>
<p>A. Recurrent episodes of binge eating (eating within a discrete period, an amount of food that is larger than what most individuals would eat in a similar period under a similar context) and a sense of lack of control over eating during the episodes</p> <p>B. The binge eating episodes are associated with three or more of the following: Eating much more rapidly than normal. Eating until feeling uncomfortably full. Eating large amounts of food when not feeling physically hungry. Eating alone because of feeling embarrassed by how much one is eating. Feeling disgusted with oneself, depressed, or very guilty afterward.</p> <p>C. Marked distress about binge eating.</p> <p>D. Binge eating occurs, on average, at least once a week for three months.</p> <p>E. binge eating is not associated with the recurrent use of inappropriate compensatory behavior as in BN and does not occur exclusively during BN or AN.</p> <p>Severity: Mild: 1-3 episodes of binge eating/per week Moderate: 4-7 Severe: 8-13 Extreme: ≥14</p>	<p>BED (of low frequency and limited duration): All of the criteria for binge-eating disorder are met, except that binge eating occurs, on average, less than once a week and for less than three months.</p>

*Adapted from American Psychological Association, "DSM-5: Eating Disorders".(pp. 350-363) [16] and [17]

2. Related works

Although eating disorders, such as anorexia nervosa, bulimia nervosa, and binge eating disorder [Table 1], have been recognized for many years, their association with Attention deficit hyperactivity disorder (ADHD) is relatively new. A better understanding was needed of the characteristics common to distinct dysregulated eating behaviors and specific moderators unique to ADHD, particularly within the pediatric population, to assist in the early identification of coexisting ADHD and eating disorders.

In 1972, Dr. Virginia Douglas, a clinical researcher at McGill University, identified a deficit in self-regulation as the prominent factor in the difficulties experienced by individuals living with Attention deficit hyperactivity disorder (ADHD). Self-regulation requires self-awareness (acknowledgment of what is occurring in the present situation), inhibition (ability

to restrain from acting on impulse and to evaluate the circumstances, options, and consequences), and both immediate and long-term strategies to attain the appropriate goal [18]. Self-regulation deficits, including deficits in emotion regulation and impulsivity, are similarly implicated in the maintenance of the eating disorder Bulimia Nervosa [Table 1] as individuals engage in binge eating or compensatory behaviors without considering long-term consequences [19][20]. Greater deficits in self-regulation have been associated with greater eating pathology of more frequent binge eating and use of compensatory behaviors [21]. Also, the inattention associated with ADHD is thought to contribute to a lack of awareness of hunger signals while focused on a task which can later lead to binge eating [22][23][24] and poor meal and snack planning [13][21].

As evidence accrues, Attention deficit hyperactivity disorder (ADHD) and eating disorders have been found to share a common psychobiological etiology of poor inhibitory control and genetic variants of the serotonergic system [26][27]. Primarily, the glucose, dopamine, and serotonin pathways and genetic receptor deficiency that influence dysregulation in executive function [27] create difficulties in inhibition, planning, and satiety [28][29]. Executive function involves neurocognitive processes that maintain appropriate problem-solving skills to attain a later goal [13][26]. Such processes are negatively impacted in individuals living with ADHD due to underdeveloped executive function [30]. The most consistent executive function deficits associated with ADHD are response inhibition, vigilance, working memory, and planning. These deficits are further believed to play a significant role in dysregulated eating behaviors [27] and influence the ability to adopt a plan and strategy to respond in a manner to attain the appropriate goal. Response processes are interrelated, as shown in imaging studies showing structural and functional alterations in the frontal and striatal brain circuits involving dopamine, primarily a deficit in the number of receptors for both ADHD and eating disorders [6]. Dopamine plays a vital role in reward and movement regulation in the brain, giving a sense of pleasure and motivation to meet essential needs. Because of this, it is believed that self-medicating with food activates the underactive dopamine pathway, increasing the level of dopamine. The inability to delay gratification (inhibitory control) and regulate arousal and focus (vigilance) impacts working memory, which guides actions of recall of consequences and the ability to plan [31]. Children with lower inhibitory control have been found to self-serve larger portions and multiple food servings [32]. As shown in [Table 1], associated impulsivity contributes to binge eating. These behaviors are considered forms of self-soothing and self-medicating with food to handle daily frustrations experienced while living with ADHD. The result is dysregulated eating patterns, and both ADHD and dysregulated eating have been found to increase the risk of obesity [15][17][33].

Literature concerning dysregulation regarding eating behaviors in the pediatric population was scarce even though binge eating [Table 1] and loss of control eating behaviors begin at a much earlier age than the diagnosis of an eating disorder [2][6][30]. Terms such as *disturbed* [34], *loss of control* [28][29], or *aberrant* [35] have been used interchangeably with the term *dysregulation* to describe abnormal eating patterns. Dysregulated eating behaviors and patterns associated with Attention Deficit Hyperactivity Disorder (ADHD) were often not clinically evaluated until the child presented with pathologic changes in appearance, body mass index, and somatic complaints requiring medical attention. Understanding characteristics associated with ADHD and dysregulated eating is the first step in changing the trajectory of dysregulated eating behaviors to reduce or prevent the development of obesity and eating disorders. Such knowledge can help improve self-esteem, self-regulation, and healthy Body Mass Index (BMI) and to avoid diabetes.

3. Method

A systematic literature review was conducted using the seven-step method of Fineout-Overholt, Melnyk, Stillwell, and Williamson [36]. This method ensured retrieval of relevant information and consisted of the following steps: Step 1: Formulating and asking a clinical question based on a practice problem. Step 2: Conducting a literature search using appropriate databases for relevant studies. Step 3: Performing a rapid critical appraisal of the studies; Step 4: Integrating the evidence with clinical expertise and patient preference and values; Step 5: Evaluating the outcomes of the practice decision based on the evidence; and Step 6: Synthesizing the evidence and presenting the results to other health providers.

3.1. Search methods

The clinical question was, "Are children living with ADHD at increased risk for dysregulated eating compared to children who do not experience ADHD? To address the question and support the implementation of a screening protocol, evidence was sought to identify patterns of eating in children diagnosed with Attention deficit hyperactivity disorder (ADHD) compared to children without ADHD. Using the PICO format, the following components were explored: population of interest (P: "children"), the concept of interest (I: "children living with ADHD"), comparison (C: "children without ADHD"), and outcome (O: "increased risk of dysregulated eating"). Synonyms for each key concept, as defined by the components of the PICO question, were explored. The second step of the method included collaboration with librarians at the university and local hospital to ensure the inclusion of appropriate databases and relevant literature.

The search entailed using PubMed, ERIC, PsycINFO, and Cumulative Index to Nursing & Allied Health Literature (CINAHL) databases with the search string "attention deficit hyperactivity disorder AND eating disorders AND children or adolescents or youth or child or teenager." Publication dates were unrestricted. Peer-reviewed journal articles published in English were included where the full text was available, and children under the age of 18 years were identified as living with Attention deficit hyperactivity disorder (ADHD). Studies incorporating children taking medications known to suppress appetite, studies of children seeking bariatric treatment, and studies involving children with comorbidities or diagnosed eating disorders were excluded. Children with defined eating disorders were excluded to ensure retrieval of research identifying eating patterns characteristic of children before fulfilling criteria for psychiatric defined eating disorders. Finally, a manual search was conducted to retrieve the citations.

3.2. Search outcome

Retrieving the studies of interest is presented in a PRISMA Flowchart [37] [Figure 1]. The retrieved studies represented research conducted in the Czech Republic, Greece, Germany, Iran, and Korea. From the initial search with Pubmed using the MeSH terms *eating*, *child*, and *attention deficit disorder with hyperactivity*, which resulted in 243 articles published between 1972 and 2020, 71 were considered for further review. Articles that did not pertain to Attention deficit hyperactivity disorder (ADHD), eating patterns in children, or eating behaviors were excluded. A concurrent search using the databases of ERIC, PsycINFO, and CINAHL was conducted using the terms *ADHD* AND "*eat* behavior*" AND *child NOT autism, adult, and medication*, resulting in 547 articles. The retrieved articles included 26 articles from PubMed and 17 from a combined database search for 44 articles. Duplicates

were removed automatically, and 44 articles that met the inclusion criteria were evaluated for relevance to the question. Due to their focus on unrelated factors (i.e., genes, animal studies, Asperger's, depression, conduct disorder, use of medication, or adults), a further 36 articles were excluded. The seven remaining articles were from PubMed.




Figure 1. Prisma search strategy pathway.

It is adapted from "Preferred Reporting Items for Systematic Reviews and Meta-Analysis: The PRISMA Statement," by D. Moher, A. Liberati, J. Tetzlaff, and D. Altman, 2010, International Journal of Surgery, vol.8, no.8, p. 658.

3.3. Quality appraisal

The next step involved critically reviewing the articles to identify key characteristics and analyzing credibility, validity, and value [38]. A quality appraisal of the studies was conducted by reviewing the research design. The studies included 4 cross-sectional studies [32][39][41], 2 case-controlled studies [42][43], and 1 experimental study [44]. The initial appraisal of the seven studies consisted of ranking each study based on the research design, from highest to lowest, as follows: Level I- Systematic review or Meta-analysis; Level II- Randomized controlled trial; Level III- Controlled trial without randomization; Level IV- Case-control or cohort study; Level V- Meta-synthesis; Level VI- Descriptive Studies; Level VII- Expert Opinion [36]. The retained studies were cross-sectional, cohort, and case-control

studies (Level IV) and experimental (Level III). Together, the studies are deemed to have moderate quality.

3.4. Design rigor

A more in-depth appraisal of the validity of the studies was conducted using a critical appraisal guide to address five essential elements (i.e., the purpose of the study, sample size, design rigor, framework, and the validity and reliability of the measurement instruments used). The purpose of all the studies was to elucidate the eating patterns of children living with Attention Deficit Hyperactivity Disorder (ADHD) and the associated characteristics that have been shown to put these children at risk of dysregulated eating and a greater tendency towards unhealthy food. The experimental study's sample size ranged from 192 to 500 children [44]. The sample sizes in the case-control studies were large enough to allow the control of confounding factors. In the cross-sectional studies using a confidence interval of 95%, the sample size allowed for a confident effect size.

Possible confounding variables such as age, sex, and socioeconomic status (SES) were identified and controlled for in 6 studies [32][39][40][42][44]. This strengthened validity. Also, validity was enhanced because the participants of most studies were not being treated with stimulants [39][40][42][43][44] or were instructed to discontinue medication for 48 hours before testing [32]. Three studies used randomization [32][39][42] [32,39,42] which minimizes selection bias. One study [41] used a convenience sample recruited through advertising at a hospital and the hospital newspaper, which could have introduced selection bias.

3.5. Framework

All seven reviewed studies used a theoretical framework or conceptual model to guide comparing eating behaviors. Six studies [32][39][40][41][42][43] used the Working Memory Model [45,46], where core deficits in ADHD (e.g., inattention, hyperactivity, and impulsivity) are theorized to impact behavioral symptoms such as dysregulated eating. The framework used in one study [44] was derived from the Tripartite Pathway Model, which attributes ADHD symptoms to deficits in inhibition, temporal processing, and motivation [47]. There is no agreement about the underlying pathway to dysregulated behaviors. However, there is agreement that the core characteristics of ADHD (e.g., impulsivity or loss of impulse control) significantly mediate loss of control eating described by Reinblatt [6] as a form of binge eating. All findings of the studies reviewed supported the premise that early problems with self-regulation are the starting point of persistent dysregulation and are characteristic of ADHD [48].

3.6. Validity of instruments

Six of the seven studies confirmed the diagnosis of ADHD by a psychiatrist [39][42] or the use of a validated culturally appropriate questionnaire [32][40][43][44]. In one study [41], an unvalidated questionnaire was translated to ensure cultural appropriateness. To evaluate eating dysregulated behaviors, four studies used validated dietary questionnaires [39][40][42][44]; only 2 [40][41] reported the instrument's reliability (Cronbach alpha as 0.77-0.89 and 0.57-0.82 respectively). All questionnaires were self-reported, which could introduce bias. Measurement error is inherent in self-reported dietary assessment instruments due to random within-person variation [49]. Individuals may overestimate portion size and

respond based on their belief of what the interviewer expects; thus, the self-report questionnaire method of gathering data has limitations. However, self-reporting was validated using interview data. Especially when considering the children's ages, this method is valid and a reasonable way of acquiring this type of information [49].

Body weight was calculated and reported in six studies using BMI and specific cut-off values for weight to height for boys and girls separately, at particular age intervals. The cut-points used in the Centers for Disease Control and Prevention (CDC) method are based on the 5th and 95th percentile BMI for age and gender. The World Health Organization (WHO) method uses cut points at two standard deviations (relative to 3rd and 97th percentiles) below and above the mean to categorize children as underweight and obese, respectively. Both these methods were used in three studies [40][41][43]. Two studies [39][42] used the International Task Force for children, while one [44] used the >90% and 97% height to weight to categorize overweight and obesity, respectively. In the studies that calculated BMI using self-reported measures of height and weight rather than direct measures (i.e., [39][42]), measurement bias was a threat to validity. One study reported significant attrition due to comorbidities, incorrect form completion, or missing data [41].

3.7. Data abstraction

Data contained in the studies that addressed findings of eating behaviors associated with Attention Deficit Hyperactivity Disorder (ADHD) were extracted. Other data extracted included descriptive information about the sample, sample size, measurement instruments, data collection, and analysis, as shown in [Table 2].

4. Results

Seven eligible studies reported on the eating behaviors of children living with Attention Deficit Hyperactivity Disorder (ADHD). Themes identified through abstraction include food quality and quantity associated with dysregulated eating behaviors. The assessment role of the nurse or other health professionals was minimally addressed in the studies.

4.1. Characteristics of study samples

As shown in [Table 2], the studies were ethnically diverse by including children from specific areas in Germany [32], Greece [41], Iran [39][42], Korea [39][43], and the Czech Republic [44]. Five studies included children between the ages of 6 and 15 [32][39][40][43][44]. One included both preschool and school-age children between the ages of 4 and 12 [42]; one focused on children who had been followed prenatally until four years of age [41]. All studies that included both sexes were found to underrepresent girls (18%-48%); only boys were included in two studies [32][44]. Leventakou [41] found that lower socioeconomic status predicted a greater prevalence of Attention Deficit Hyperactivity Disorder (ADHD).

Table 2. Selected data extracted from reviewed studies

Author	Abbasi et al. [42]	Azadbakht et al. [39]	Kim et al. [40]	Leventakou et al. [41]	Ptacek et al. [34]	Wilhelm et al. [44]	Woo et al. [43]
Year	2019	2012	2018	2016	2014	2011	2014
Country	Iran	Iran	Korea	Greece	Czech Republic	Germany	Korea
Framework	Working memory Model	Working memory Model	Working memory Model	Working memory Model	Working memory Model	Tripartite Pathway Model	Working memory Model
Design Method	Cross-sectional	Case-Control	Cross-sectional	Cross-sectional	Cross-sectional	Quasi-experiment	Case-Control
Participant age	4-12 Mean 7 +/- 1	7-9 Mean 8 ± 1	5-13 Mean 9.4+/-1.7	4 Mean 4	6-10 Mean NR	7-15 Mean NR	7-12 Mean 9± 1
Participant gender	N=500 M=82	N=375 M=52%	N=12 350 M= 49%	N= 471 M=59 %	N=200 M=100%	N=94 M=100%	N=192 M=34%
Setting	Isfahan Psych clinic	Tehrani elementary schools	Cheonan elementary schools	Create 2007 Birth cohort	Prague	Aachen Hospital	Busan 2 University Hospitals
Instrument	BMI M DSM-5 D FFQ	BMI M DSM IV Q FFQ	BMI SR DSM IV FFQ	BMI SR DSM IV Q BEF CEBQ	BMI NR DSM IV Conner's	BMI DSM IV D FBB HKS	BMI NR DSM IV D
Findings	Diet Patterns Processed Calorie dense High sugar Healthy	Diet Patterns Sweet Fast food Healthy High sugar	Behaviors High sugar Overeat > Speed	Behaviors Emotional eating Overeat Eat not hungry High food response	Behaviors Skip meals Meals >5/d Sugar drink preference	Behaviors > speed (1 st 30 min) Uncontrolled snacking Delayed aversion	Diets Tradition Seaweed-egg Low fat-high carb Healthy snack

*Note. BEF BMI M=body mass index- measured; BMI-SR=body mass index-self reported; CEBQ= child eating behavior questionnaire-parent; DSM IV Diagnostic and Statistical Manual Fourth Edition; D= Diagnosed, Q= Questionnaire; DSM-5 = fifth edition; FFB HKS = German Conners Questionnaire; FFQ= food frequency questionnaire; M= male; NR not reported;

4.2. Food quality

Of the five studies evaluating eating characteristics related to food preference, two [39][42] reported that children with Attention deficit hyperactivity disorder (ADHD) demonstrated a higher preference for high sugar, high fat, highly processed, and high caloric foods, and a pattern of high consumption of fast food compared to other children. Children with the highest *Western dietary pattern* rating, based on eating high-fat and high-calorie foods, had greater odds of having ADHD in comparison to children in the healthy eating pattern, which was inversely associated with ADHD (OR = 0.46; 95% CI, 0.38-0.91; $p=0.01$) [42]. In evaluating patterns of eating different types of food, children with ADHD were reported as being at greater risk of choosing and consuming foods and snacks categorized as *unhealthy* [40][44]. The foods categorized as *unhealthy*, high in sugar and fat, were processed and included fast foods, soft drinks, and instant noodles. Sugary drinks make up almost half of the children's daily intake [44]. Alternatively, children without ADHD were reported to have a traditional healthy diet [43].

4.3. Food quantity

Children living with Attention deficit hyperactivity disorder (ADHD) were found to exhibit markedly diminished adherence to a traditional breakfast, lunch, and dinner schedule and significantly higher frequency (>5/day) of irregular eating times [32]. A significant association between ADHD and overeating behavior was associated with the consumption of *unhealthy* food, the number of overeating episodes per week, and eating speed [44].

4.4. ADHD and eating behaviors

The majority of outcomes reported a significant effect of ADHD on eating behaviors in children [32][39][40][41][42][44]; one study found an association with healthy traditional eating predictive of lower incidences of ADHD [43]. In all the reviewed studies, researchers focussed on impulsivity, inattentiveness, and hyperactivity as the core characteristics of Attention Deficit Hyperactivity Disorder (ADHD) that mediate the relationship between ADHD and dysregulated eating behaviors. Increased ADHD symptoms were associated with food responsiveness (reward sensitivity) and emotional overeating (eating to temper negative emotions) in preschool children [41]. This suggests that impulsivity may play an essential role in creating vulnerability to binge eating [6]. Furthermore, a lack of impulse control, an inhibitory deficit, and persistent difficulties typical in ADHD (e.g., inattention to/diminished awareness of hunger and satiety cues, distractibility, and hyperfocus on ingesting sweet or salty foods that stimulate dopamine) increase the likelihood of episodes of overeating and eating in the absence of hunger. This result is supported by Yao et al. [50], who found the prevalence of ADHD symptoms as high as 35% to 37% for BN and AN binge eating/purging subtype versus 18% for AN restricting subtype in a large Swedish sample (N =1165) of adults clinically diagnosed as having eating disorders. Yao et al. additionally found that common genetic risk variants for ADHD were significantly associated with eating disorder symptoms and that ADHD symptoms during early childhood predicted binge-eating behaviors during adolescence. These results help demonstrate why eating disorders and ADHD co-occur and underscore the importance of screening for eating disorder symptoms in individuals who experience ADHD.

5. Discussion

The systematic review provided a summary of the results of seven studies that examined the eating behaviors of children living with Attention Deficit Hyperactivity Disorder (ADHD). The core ADHD characteristics of impulsivity, inattention, and hyperactivity were found to be associated with dysregulated eating behaviors. ADHD affects eating behavior due to the difficulties affecting the ability to plan, regulate the desire for rewards, and manage daily negative experiences. These difficulties increase eating irregularly, eating to feel better, and, in turn, overeating.

In the studies reviewed, the majority of outcomes evaluated were related to food preference, with only two studies evaluating the schedule of eating, which could significantly influence the risk of obesity [32][40]. One study found low socioeconomic status predicted a greater prevalence of Attention deficit hyperactivity disorder (ADHD) [41]. This finding is worth further consideration as Biederman et al. [48] and Russell et al. [49] similarly found a higher prevalence of ADHD with lower socioeconomic status. In general, low socioeconomic status is reported to increase the likelihood of poor mental health in children by 1.18 to 3.34 times [49] and specifically relates to a higher prevalence of ADHD [48][49][60]. The

underlying cause has yet to be elucidated; however, it is understood that food availability is influenced socially, culturally, economically, and geographically.

In the studies, sample sizes were appropriate but were too small to further differentiate ADHD by their subtypes, which are categorized by prominent symptoms. Many studies identified children of the combined subtype (inattention and hyperactivity) as the predominant subtype in males. Leventakou and colleagues' (2016) findings were in contrast to previous studies in which executive function deficits were found to mediate dysregulated eating [26][29][57]. Executive function is related to self-regulation, a skill developed within the first decade of life that regulates behavior, cognition, and emotion. The contrasting results may have been due to the fit of the measurement tool with the age of the children. The children were four years of age, and the evaluation tool used may not have been adequate for evaluating executive function in this age group. Attentional control through the regulation of cognition and emotion has been found to develop substantially between the ages of three and four years, with further substantial effortful control through age seven [58]. Therefore, an appreciable difference may have yet to be identified at this age. Future research should seek clarification as to whether the subtypes of ADHD, categorized by symptoms, significantly predict eating behaviors. This research would need larger sample sizes, consistently validated measurement instruments, and direct height, weight, and intake measures. Professional practice and academic programs could provide a more comprehensive curriculum on children's mental health to equip healthcare providers better to address dysregulated eating behaviors in healthcare settings effectively.

5.1. ADHD, food, and eating behaviors

The findings of high sugary, high-carbohydrate food ingestion reported in some studies of this review [34][39][40][42] may indicate seeking out foods that increase serotonin and dopamine, providing a sense of pleasure and calmness [53][54][55]. Serotonin contributes to feelings of well-being, reduced irritability, and improved overall mood and learning; dopamine plays a critical role in the reward pathway [23]. Positive emotional eating activates the dopaminergic system, initiating behavior to obtain the reward and reinforcing the consumption of certain foods [57]. People whose brains are low in dopamine often self-soothe with food because of its ability to temporarily activate dopamine. For both individuals living with Attention Deficit Hyperactivity Disorder (ADHD) and those with obesity or eating disorders, the brain is slower to absorb glucose [51]. This reduced glucose metabolism, together with decreased receptors for dopamine, makes less energy available to the attention center (in the prefrontal cortex) that activates executive function (such as focus and sustained attention) [54][55]. Subsequently, increased Attention Deficit Hyperactivity Disorder (ADHD) symptoms are associated with food responsiveness (reward sensitivity) and emotional overeating (eating to temper negative emotions) in preschool children [41]. Higher sensitivity to reward has been associated with a stronger enjoyment of food, higher food responsiveness (i.e., desire to eat in the absence of hunger or satiety), and more emotional overeating when experiencing negative emotions [50]. Individuals who experience ADHD and bulimia nervosa have decreased dopamine receptor binding and release [54][56]. Imaging studies have found structural and functional alterations in the frontal and striatal regions in both ADHD and eating disorders, notably altered brain circuits involving dopamine [56]. The reward deficiency theory is closely linked as it is believed that self-soothing with food activates the underactive dopamine pathway, resulting in increased levels of dopamine [51][52]. This is a shared process for both individuals with ADHD and binge eating disorder

and a maladaptive way of regulating dopamine levels [56]. It is important to note, however, that even though children with ADHD prefer high fat, high sugar, and calorically dense food and eat more significant amounts more frequently, younger children may present with average weight or underweight [59]. This emphasizes the critical need for further evaluation in this age group.

5.2. Implications for nurse practitioners

Many Nurse Practitioners (NPs) work in primary care settings, making these NPs well-positioned to inquire into and evaluate eating patterns during scheduled well-child visits. The NP performs patient-directed health assessments and diagnoses and manages acute and chronic conditions [61]. NPs and other healthcare providers use a standard evidence-based infant/child health maintenance tool (the Rourke) to evaluate growth and development during routine child wellness visits [62]. However, this tool does not guide how to assess eating behaviors. The Canadian Practice Guidelines for ADHD have been developed and reviewed by a multidisciplinary team of Attention deficit hyperactivity disorder (ADHD) specialists, pediatricians, psychiatrists, psychologists, family physicians, nurses, educators, and community stakeholders from Canada and the US. The guidelines are accessible through the Canadian ADHD Resource Alliance [63] and guide diagnosis, management, and treatment across the lifespan for individuals with ADHD. The recent identification of the associated risk of eating disorders and obesity related to dysregulated eating is discussed; however, guidance or resources on how to approach the issue with family and individuals are not provided. To effectively enact the health care provider role, awareness of the manifestations of dysregulated eating will ensure comprehensive inquiry about dietary patterns and the use of appropriate resources.

The synthesized findings of this review highlight the need for supplemental evaluation of eating patterns for children living with dysregulated eating of ADHD due to the risk of inadequate nutrition, obesity, and eating disorder development. The patterns between ADHD and eating disorders suggest that family history may help with early detection and risk identification of the two disorders. ADHD has been shown to predict eating disorder symptoms; regular screening for ADHD symptoms in those with eating disorders and for eating disorder symptoms in those with ADHD may hasten the detection of and appropriate intervention for comorbid conditions [50]. Thus, this study's findings alert healthcare providers of the necessity to screen for both eating disorders and ADHD when symptoms appear for either disorder. When interviewing and counseling pediatric patients and their parents regarding eating behaviors in the context of ADHD, understanding the manifestations and implications of dysregulated eating patterns is necessary. Individual barriers to disclosing non-normative eating include associated stigma and awareness of societal prejudices and discrimination against those with ADHD [55]. Providing education to the general population can assist people in identifying symptoms or risk behaviors, lessening the stigma, and seeking early advice to improve outcomes.

5.2. Limitations

Given that all the studies were derived from medical research and the role of the nurse practitioner and nursing care strategies were not addressed, this area requires further research. As well, the studies were predominantly focused on male children, and this contributes to unrecognized ADHD issues in females before adolescence when eating disorders become more prevalent. Although ADHD and obesity are global issues, the generalizability of our

synthesis is limited due to cultural variability (i.e., there were no studies representing children from Canada or the United States). However, the seven included studies reported similar prevalence of ADHD, and, even though cultural food differences were identified, all the studies consistently found significantly higher intakes of fast food, sugary foods and beverages, and calorically dense foods with children diagnosed with ADHD. These behaviors are detrimental to physiological and psychological development and health.

6. Conclusion

This study was conducted to identify features of dysregulated eating in the context of children living with Attention Deficit Hyperactivity Disorder (ADHD). The overall methodological quality of the studies was moderate. The evidence suggests underlying core characteristics of ADHD, such as impulsivity, inattention, and emotional dysregulation, are mediators in dysregulated eating behaviors. An awareness of how executive function dysregulation impacts psychological, behavioral, cognitive, and social domains provides a better understanding of the associated problems and complexity of eating behaviors related to ADHD in children. Nurses and other health professionals can encourage open dialogue about the daily lived experiences of individuals with ADHD and, in particular, identify dysregulated eating. However, it is not known if all children with ADHD have dysregulated eating or which children will develop eating disorders.

Further exploration is needed to determine identifiable characteristics of dysregulated eating to facilitate early recognition and intervention. This review encourages further rigorous research studies to provide stronger evidence of the different manifestations of disordered eating in the context of ADHD. There is a need for high-quality research that provides more descriptive evidence of dysregulated eating in the various subtypes of ADHD, and that includes girls. Given the disproportionate rate of eating disorders, obesity, depression, and other challenges among children living with ADHD, early identification of dysregulated eating behaviors is crucial.

Acknowledgments

The authors thank Paul Clark and Richelle Witherspoon for conducting the initial search to ensure the search strategy was comprehensive.

Conflict of interest

The authors have declared no conflict of interest.

References

- [1] K. R. Krull, "Attention deficit hyperactivity disorder in children and adolescents: Clinical features and diagnosis," UpToDate, (2022) <https://www.uptodate.com/contents/attention-deficit-hyperactivity-disorder-in-children-and-adolescents-clinical-features-and-diagnosis>
- [2] R. Barkley, "Attention-deficit hyperactivity disorder," 4th ed. Guilford Publications, New York, (2018)
- [3] S. Sulkes, "Attention-deficit/hyperactivity disorder (ADD, ADHD)," Merck Manuals Professional Edition, (2018), <https://www.merckmanuals.com/professional/pediatrics/learning-and-developmental-disorders/attention-deficit-hyperactivity-disorder-add,-adhd>

- [4] W. B. Daviss, "A review of co-morbid depression in pediatric ADHD: etiology, phenomenology, and treatment." *Journal of Child Adolescent Psychopharmacology*, vol.18, no.6, pp.565-571, (2008) DOI:10.1089/cap.2008.032
- [5] J. Mitchell, C. Robertson, A. Anastopoulos, R. Nelson-Gray, and S. Collins, "Emotion dysregulation and emotional impulsivity among adults with attention-deficit/hyperactivity disorder: Results of a preliminary study," *Journal of Psychopathology and Behavioral Assessment*, vol.34, no.4, pp.510-519, (2012) DOI:10.1007/s10862-012-9297-2
- [6] S. Reinblatt, L. Jeannie-Marie, M. Mahone, S. Forrester, H. Wilcox, and M. Riddle, "Association between binge eating and attention-deficit/hyperactivity disorder in two pediatric community mental health clinics," *International Journal of Eating Disorders*, vol.48, no.5, pp.505-511, (2015) DOI:10.1002/eat.22342
- [7] T. Hanć, and S. Cortese, "Attention-deficit/hyperactivity disorder and obesity: A review and model of current hypotheses explaining their comorbidity," *Neuroscience Biobehavioural Review*, vol.92, pp.16–28. 2018 DOI:10.1016/j.neubiorev.2018.05.017
- [8] M. Waring and K. Lapane, "Overweight in children and adolescents in relation to attention-deficit/hyperactivity disorder: Results from a national sample," *Pediatrics*, vol.122, no.1, pp.1-6, (2008) DOI:10.1542/peds.2007-1955
- [9] S. Donnchadha, J. Bramham, and C. Greene, "Rethinking the association between overweight/obesity and ADHD in children: a longitudinal and psychosocial perspective," *Irish Journal of Psychological Medicine*, Jan 24, pp.1-14, (2020) DOI:10.1017/ipm.2019.61
- [10] J. Bleck, R. DeBate and R. Olivardia, "The comorbidity of ADHD and eating disorders in a nationally representative sample," *The Journal of Behavioral Health Services & Research: Official Publication of the National Council for Behavioral Health*, vol.42, no.4, pp.437-451, (2015) DOI: 10.1007/s11414-014-9422-y
- [11] A. Agranat-Meged, C. Deitcher, G. Goldzweig, L. Leibenson, M. Stein and E. Galili-Weisstub, "Childhood obesity and attention-deficit/hyperactivity disorder: A newly described comorbidity in obese hospitalized children," *International Journal of Eating Disorders*, vol.37, no.4, pp.357-359, (2005) DOI:10.1002/eat.20096
- [12] R. Strauss, "Childhood obesity and self-esteem," *Pediatrics*, vol.105, no.1, p.513, (2000) DOI:10.1097/00005176-199910000-00124
- [13] B. P. Nazar, C. Bernardes, G. Peachey, J. Sergeant, P. Mattos, and J. Treasure, "The risk of eating disorders comorbid with attention-deficit/hyperactivity disorder: A systematic review and meta-analysis," *International Journal of Eating Disorders*, vol.49, no.12, pp.1045-1057, (2016) DOI:10.1002/eat.22643
- [14] B. Fuemmeler, T. Ostbye, C. Yang, J. McClermon, and S. Kollins, "Association between attention-deficit/hyperactivity disorder symptoms and obesity and hypertension in early adulthood: A population-based study," *International Journal of Obesity*, vol.35, no.6, pp.852–862, (2011) DOI:10.1038/ijo.2010.214
- [15] J. T. Nigg, J. M. Johnstone, E. D. Musser, H .G. Long, M. T. Willoughby and J. Shannon, "Attention-deficit/hyperactivity disorder (ADHD) and being overweight/obesity: New data and meta-analysis," *Clinical Psychology Review*, vol.43, pp.67-79, (2016) DOI:10.1016/j.cpr.2015.11.005
- [16] American Psychiatric Association, "Binge eating disorder," *Diagnostic and statistical manual of mental disorders (5th ed.)*. Author, Arlington, VA, (2013)
- [17] C. Curtin, L. Bandini, E. Perrin, D. Tybor, and A. Must, "Prevalence of overweight in children and adolescents with attention deficit hyperactivity disorder and autism spectrum disorders: A chart review," *BioMed Central Pediatrics*, vol.5, no.1, p. 48, (2005) DOI:10.1186/1471-2431-5-48
- [18] V. Douglas, "Cognitive deficits in children with attentional deficit hyperactivity disorder. A long-term follow-up," *Canadian Psychology*, vol.46, no.1, pp.23-31, (2005) DOI:10.1037/h0085821
- [19] E. K. Presseller, M. L. Wilkinson, C. Trainor, E. W. Lampe, and A. S. Juarascio, "Self-regulation deficits moderate treatment outcomes in a clinical trial evaluating just-in-time adaptive interventions as an augmentation to cognitive-behavioral therapy for bulimia-spectrum eating disorders," *International Journal of Eating Disorders*, vol.55, no.5, pp.709-716, (2022) DOI:10.1002/eat.23695. Epub 2022 Feb 24

- [20] N. E. Svedlund, C. Norring, Y. Ginsberg, and Y. von Hausswolff-Juhlin, "Symptoms of attention deficit hyperactivity disorder (ADHD) among adult eating disorder patients," *BMC Psychiatry*, vol.17, no.19, (2017) DOI:10.1186/s12888-016-1093-1
- [21] K. E. Smith, T. B. Mason, L. M. Schaefer, A. Juarascio, R. Dvorak, N. Weinbach, R. D. Crosby, and S.A. Wonderlich, "Examining intra-individual variability in food-related inhibitory control and negative affect as predictors of binge eating using ecological momentary assessment," *Journal of Psychiatric Research*, vol.120, Jan, pp.137-143, (2020) DOI:10.1016/j.jpsychires.2019.10.017
- [22] J. Fleming and L. Levy, "ADHD and disordered eating," In P. Quinn and K. Nadeau (Eds.), *Gender issues and AD/HD: Research, diagnosis, and treatment*, Silver Spring, MD: Advantage Books, pp. 370-382 (2002)
- [23] C. Davis, R. Levitan, M. Smith, S. Tweed, and C. Curtis, "Associations among overeating, overweight, and attention-deficit/hyperactivity disorder: A structural equation modeling approach," *Eating Behaviors*, vol.7, no.3, pp.266-274, (2006) DOI: 10.1016/j.eatbeh.2005.09.006
- [24] U. Pauli-Pott, B. Becker, O. Albayrak, J. Hebebrand, and W. Pott, "Links between psychopathological symptoms and disordered eating behaviors in overweight/obese youths," *International Journal of Eating Disorders*, vol. 46, no.2, pp.156-163, (2013) DOI:10.1002/eat.22055
- [25] Y. Lee, and P. Lin "Association between serotonin transporter gene polymorphism and eating disorders: A meta-analytic study," *International Journal of Eating Disorders*, vol.43, no.6, pp.498-504, (2010) DOI:10.1002/eat.20732
- [26] N. Stulz, U. Hepp, C. Gächter, C. Martin-Soelch, A. Spindler, and G. Milos, "The severity of ADHD and eating disorder symptoms: A correlational study," *BioMed Central Psychiatry*, vol.13, no.1, pp.44-50, (2013) DOI:10.1186/1471-244X-13-44
- [27] M. Semrud-Clikeman, J. Walkowiak, A. Wilkinson, and B. Butcher, "Executive functioning in children with Asperger syndrome, ADHD-combined type, ADHD-predominantly inattentive type, and controls." *Journal of Autism and Developmental Disorders*, vol.40, no.8, pp.1017-1027, (2010) DOI: 10.1007/S10803-010-0951-9
- [28] A. Hilbert, S. Kurz, D. Dremmel, S. Blüher, S. Munsch and R. Schmidt, "Cue reactivity, habituation, and eating in the absence of hunger in children with loss of control eating and attention-deficit/hyperactivity disorder," *International Journal of Eating Disorders*, vol.51, no.3, pp.223-232, (2018) DOI:10.1002/eat.22821
- [29] S. Kurz, D. Schoebi, D. Dremmel, W. Kiess, S. Munsch, and A. Hilbert, "Satiety regulation in children with loss of control eating and attention-deficit/hyperactivity disorder: A test meal study," *Appetite*, vol.116, pp. 90-98 (2018). DOI: 10.1016/j.appet.2017.04.013
- [30] Z. Yilmaz, N. Kristin, K. Javaras, J. Baker, L. Thornton, P. Lichtenstein, C. M. Bulik, and H. Larsson, "Association between childhood to adolescent attention-deficit/hyperactivity disorder symptom trajectories and late adolescent disordered eating," *Journal of Adolescent Health*, vol.61, no.2, pp.140-146, (2017) DOI:10.1016/j.jadohealth.2017.04.001
- [31] J. Holmes, K. Hilton, M. Place, T. Alloway, J. Elliott, and S. Gathercole, "Children with low working memory and children with ADHD: Same or different?" *Frontiers in Human Neuroscience*, vol.8, no.976, pp.1-13, (2014) DOI:10.3389/fnhum.2014.00976
- [32] A. Fogel, K. McCrickerd, A. Goh, L. Fries, Y. Chong, K. Tan, F. Yap, L. P. Shek, M. J. Meaney, S. Cai, P. Pelufo Silveira, B. F. P. Broekman, Y. S. Lee, K. M. Godfrey, M. F. F. Chong, and C. Forde, "Association between inhibitory control, eating behaviors and adiposity in 6-year-old children," *International Journal of Obesity*, vol.43, no.7, pp.1344-1353 (2019) DOI:10.1038/s41366-019-0343-y
- [33] M. Erhart, B. Herpertz-Dahlmann, N. Wille, B. Sawitzky-Rose, H. Hölling, and U. Ravens-Sieberer, "Examining the relationship between attention deficit/-hyperactivity disorder and overweight in children and adolescents," *European Child & Adolescent Psychiatry*, vol.21, no.1, pp.39-49. (2012) DOI:10.100/s00787-011-0230-0
- [34] R.Ptacek, H. Kuzelova, G. Stefano, J. Raboch, T. Sadkova, M. Goetz, and R.Kream, "Disruptive patterns of eating behaviors and associated lifestyles in males with ADHD," *Medical Science Monitor: International*

- Medical Journal of Experimental and Clinical Research, vol.20, pp. 608-613, (2014)
DOI:10.12659/MSM.890495
- [35] S. Cortese, Commentary: Switching the zoom on the ADHD research lens – A reflection on Leventakou et al.," *Journal of Child Psychology and Psychiatry*, vol.57, no.6, pp. 685–686 (2016) DOI:10.1111/jcpp.12555
- [36] E. Fineout-Overholt, B. Melnyk, S. Stillwell, and K. Williamson, "Critical appraisal of the evidence: Part III. The process of synthesis: Seeing similarities and differences across the body of evidence," *American Journal of Nursing*, vol.110, no.11, pp. 43-51, (2010) DOI:10.1097/01NAJ.0000390523.99066.b5
- [37] D. Moher, A. Liberati, J. Tetzlaff, and D. Altman, "Preferred reporting items for systematic reviews and meta-analysis: The PRISMA statement," *International Journal of Surgery*, vol.8, no.5, pp. 336-41, (2010) DOI:10.1016/j.jisu.2010.02.007. Epub 2010 Feb 18. Erratum in: *Int J Surg*. 2010, vol.8, no.8, pp.658. PMID: 20171303, (2010)
- [38] K. Scott and R. McSherry, "Evidence-based nursing: Clarifying the concepts for nurses in practice," *Journal of Clinical Nursing*, vol. 18, no.11, pp. 1085-1095, (2009) DOI:10.1111/j.1365-2702.2008.02588.x
- [39] L. Azadbakht and A. Esmailzadeh, "Dietary patterns and attention deficit hyperactivity disorder among Iranian children," *Nutrition*, vol.28, no.3, pp.242-249, (2012). DOI: 10.1016/j.nut.2011.05.018
- [40] K. M. pt, M. H. Lim, H.-J. Kwon, H., S.-J. Yoo, E.-J. Kim, J. W. Kim, M. Ha, and K. C. Paik, "Associations between attention-deficit/hyperactivity disorder symptoms and dietary habits in elementary school children," *Appetite*, vol.127, pp. 274-279, (2018) DOI:10.1016/j.appet.2018.05.004
- [41] V. Leventakou, I. N. Micali, V. Georgiou, K. Sarri, I. K. Koutra, S. Koinaki, I. M. Vassilaki, I. M. Kogevinas, and L. Chatzi, "Is there an association between eating behavior and attention-deficit hyperactivity disorder symptoms in preschool children?" *Journal of Child Psychology and Psychiatry*, vol.57, no.6, pp.676-684, (2016) DOI:10.1111/jcpp.12504
- [42] K. Abbasi, S. Beigrezai, R. Ghiasvand, M. Pourmasoumi, and B. Mahaki, "Dietary patterns and attention deficit hyperactivity disorder among Iranian children: a case-control study," *Journal of the American College of Nutrition*, vol.38, no.1, pp. 76-83. (2019). <https://doi.org/10.1080/07315724.2018.1473819>
- [43] H. D. Woo, D. W. Kim, Y.-S. Hong, Y.-M. Kim, J.-H. Seo, B. M. Choe, J. H. Park, J.-W. Kang, J.-H. Yoo, H. W. Chueh, J. H. Lee, M. J. Kwak, and J. Kim. "Dietary patterns in children with attention-deficit/hyperactivity disorder (ADHD)," *Nutrients*, vol.6, pp.1539-1553, (2014) DOI: 10.3390/nu6041539
- [44] C. Wilhelm, I. Marx, K. Konrad, K. Willmes, K. Holtkamp, T. Vloet, and B. Herpertz-Dahlmann, "Differential patterns of disordered eating in subjects with ADHD and overweight," *The World Journal of Biological Psychiatry*, vol.12, sup1, pp.118-123, (2011) DOI:10.3109/15622975.2011.602225
- [45] A. D. Baddeley and G. Hitch, "Working memory," In G. H. Bower (Ed.), *The Psychology of learning and motivation: Advances in research and theory*, Academic Press New York, pp.47-89 (1974) DOI:10.1016/S0079-7421(08)60452-1
- [46] M. Rapport, K. Chung, G. Shore, and P. Isaacs, "A conceptual model of child psychopathology: Implications for understanding attention deficit hyperactivity disorder and treatment efficacy," *Journal of Clinical Child Psychology*, vol.30, no.1, pp. 48-58, (2001) DOI:10.1207/S15374424JCCP3001_6
- [47] E. Sonuga-Barke, P. Bitsako, and M. Thompson, "Beyond the dual pathway model: Evidence for the dissociation of timing, inhibitory, and delay-related impairments in attention-deficit/hyperactivity disorder," *Journal of the American Academy of Child & Adolescent Psychiatry*, vol.49, no.4, pp. 345-355, (2010) DOI:10.1176/appi.ajp.2012.12070991
- [48] J. Biederman, E. Mick, S. Faraone, E. Braaten, A. Doyle, T. Spencer, T. Wilens, E. Frazier, and M. Johnson, "Influence of gender on attention deficit hyperactivity disorder in children referred to a psychiatric clinic," *American Journal of Psychiatry*, vol.159, no.1, pp.36-42, (2002) DOI: 10.1176/appi.ajp.159.1.36
- [49] A. Russell, T. Ford, R. Williams, and G. Russell, "The Association Between Socioeconomic Disadvantage and Attention Deficit/Hyperactivity Disorder (ADHD): A Systematic Review," *Child Psychiatry Human Development*, vol.47, no.3, pp.440-458, (2016) DOI:10.1007/s10578-015-0578-3

- [50] S. Yao, R. Kuja-Halkola, J. Martin, Y. Lu, P. Lichtenstein, C. Norring, A. Birgegård, Z. Yilmaz, C. Hübel, H. Watson, J. Baker, and C. Almqvist, Eating Disorders Working Group of the Psychiatric Genomics Consortium, L. M. Thornton, P. K. Magnusson, C. M. Bulik, and H. Larsson, "Associations between attention-deficit/hyperactivity disorder and various eating disorders: A Swedish nationwide population study using multiple genetically informative approaches," *Biol Psychiatry*, vol.86, no.8, pp.577-586 (2019) DOI 10.1016/j.biopsych.2019.04.036
- [51] M. Michaelides, J. Pascau, J-D. Gispert, F. Delis, D. Grandy, G-J. Wang, M. Desco, M. Rubinstein, N.D. Volkow, and P. Thanos, "Dopamine D4 receptors modulate brain metabolic activity in the prefrontal cortex and cerebellum at rest and in response to methylphenidate" (2010) DOI:10.1111/j.1460-9568.2010.07319
- [52] R. Johnson, M. Gold, D. Johnson, T. Ishimoto, M. Lanaspa, N. Zahniser, and N. Avena, "Attention-Deficit/Hyperactivity Disorder: Is it time to Reappraise the Role of Sugar Consumption?" *Postgraduate Medicine*, vol.123, no.5, pp.39-49, (2011), DOI:10.3810/pgm.2011.09.2458
- [53] A. Mueller, D. Hong, S. Shepard, and T. Moore, "Linking ADHD to the Neural Circuitry of Attention. *Trends Cognitive Science*, vol.21, no.6, pp.474-488, (2017) DOI: 10.1016/j.tics.2017.03.009
- [54] N. D. Volkow, G. J. Wang, F. Telang, J. S. Fowler, P. K. Thanos, J. Logan, D. Alexoff, Y. S. Ding, C. Wong, Y. Ma, and K. Pradhan, "Low dopamine striatal D2 receptors are associated with prefrontal metabolism in obese subjects: possible contributing factors," *Neuroimage* vol.42, no.4, pp. 1537-1543, (2008) DOI:10.1016/j.neuroimage.2008.06.002
- [55] N. D. Volkow, G. F. Wang, S. H. Kollins, T. L. Wigal, J. H. Newcorn, F. Telang, J.S. Fowler, W. Zhu, J. Logan, Y. Ma, K. Pradhan, C. Wong, and J.M. Swanson, "Evaluating dopamine reward pathway in ADHD: clinical implications," *JAMA*, vol.302, no.10, pp.1084-1091, (2009) DOI:10.1001/jama.2009.1308. Erratum in: *JAMA*. 2009 Oct 7, vol.302, no.13, pp.1420
- [56] C. Davis, K. Patte, R.D. Levitan, J. Carter, A.S. Kaplan, C. Zai, C. Reid, C. Curtis, and J. L. Kennedy, "A psycho-genetic study of associations between the symptoms of binge eating disorder and those of attention deficit (hyperactivity) disorder," *Journal of Psychiatric Research*, vol.43, no.7, pp. 687-696, (2009) DOI:10.1016/j.jpsychires.2008.10.010
- [57] A. Goldschmidt, S. O'Brien, J. Lavender, C. Pearson, D. LeGrange, D., and S. Hunter, "Executive functioning in a racially diverse sample of children who are overweight and at risk for eating disorders," *Appetite*, vol. 124, pp.43-49. (2018) DOI: 10.1016/j.appet.2017.03.010
- [58] E. J Kim, H. J. Kwon, M. Ha, M. H. Lim, S.Y. Oh, J. H. Kim, S. J. Yoo, and K. C. Paik "ADHD, dietary behaviours and obesity," *Child Care Health Development*, vol.40, pp. 698-705, (2014) DOI:10.1111/cch.12129
- [59] T. Hanć, A. Słopeń, T. Wolańczyk, A. Szwed, Z. Czapla, M. Durda, M. Dmitrzak-Węglarz, and J. Ratajczak. "Attention-Deficit/Hyperactivity Disorder is Related to Decreased Weight in the Preschool Period and Increased Rate of Overweight in School-Age Boys," *Journal of Child and Adolescent Psychopharmacology*, vol.25, no.9, pp.691-700, (2015) DOI:10.1089/cap.2014.0157
- [60] A. Russell, T. Ford, and G. Russell, "Socioeconomic associations with ADHD: Findings from a mediation analysis," *PloS one*, vol. 10, no. 6, e0128248, (2015) DOI:10.1371/journal.pe.0128248
- [61] Canadian Nurses Association, "Position statement on mental health services," (2012) https://www.cna-aic.ca/-/media/can/page-content/pdf-en/ps85_mental_health_e.pdf?la=en.2012
- [62] L. Rourke, D. Leduc, and J. Rourke, "Rourke baby record: Evidence-based infant/child health maintenance." (2017) www.rourkebabyrecord.ca
- [63] Canadian Attention Deficit Hyperactivity Disorder Resource Alliance, "Canadian ADHD Practice Guidelines," (3rd ed.) CADDRA, Toronto ON (2018) <https://www.caddra.ca>
- [64] A. Mueller, A. Fermaier, J. Koerts, and L. Tucha, "Stigma in attention deficit hyperactivity disorder," *Attention Deficit Hyperactivity Disorder*, vol.4, no.3, pp.101-114 (2012) DOI:10.1007/s12402-012-0085-3

Authors



Janice Arsenault

Primary Care Nurse Practitioner, ADHD and Beyond.



Kathryn Weaver

Honorary Research Professor, University of New Brunswick